

# Long-Term Murine Keratinocyte Cultures Become Tetraploid, Yet Maintain the Ability to Stratify

To the Editor:

We would like to convey a word of caution to readers of the Journal. Two recent reports in this journal describe methods for long-term cultivation of newborn mouse keratinocytes (Hager *et al*, 1999; Caldelari *et al*, 2000). Both reports strongly implied, without disclaimers or caution, that normal mouse keratinocytes could be grown long term from single mouse pups. Hager *et al*, introduce their paper by noting the desirability of culturing mouse keratinocytes from single pups, especially by those who might have generated mice with characterized mutant genes. Indeed, we undertook the use of the Hager method for that purpose and were surprised to find that the cells became polyploid after as few as five passages.

We easily established cell lines from single newborn pups, representing eight different mouse strains and genetically defined mutants. Cells were subcultured every 7–10 d and became homogeneous in appearance and in growth rate, usually after 2 mo in culture or after five to seven passages. After eight to ten passages, five cell lines were tested and found to be nearly tetraploid, as determined by chromosome counts on at least 20 spread mitotic cells (Table I).


All of our lines have the expected characteristics of normal differentiation described by Hager *et al*, and Caldelari *et al*: an epithelial, cobblestone appearance at confluence and expression of biochemical markers of differentiation following elevation of the medium calcium concentration (for example, induction of K1; Fig 1A). Moreover, we found that late-passage keratinocytes retain the ability to form a stratified, if somewhat disorganized, epithelium (Fig 1B). Some even form a stratum corneum and express late markers of epidermal differentiation, such as keratohyalin granules (Fig 1Bb) and immunohistochemical evidence of loricrin expression (not shown).

**Table I. Chromosome Analysis of Cultured Mouse Keratinocytes**

Strain (transgene)	Chromosomes (weeks <i>in vitro</i> at analysis)	Weeks <i>in vitro</i> to date
BALB/c	80 ± 2.3 (14) <sup>a</sup>	26
C57BL/6 (SupFG1)	70 ± 16 (9)	33
C57BL/6J ( <i>hprt</i> <sup>-/-</sup> )	80 ± 3.3 (13)	21
129/SvEvTac ( <i>hfe</i> <sup>-/-</sup> )	78 ± 4.8 (22)	24
SKH1	77 ± 0.5 (5)	16

<sup>a</sup>Mean ± SD.

It is well known that cultured murine cells are prone to chromosomal instability. With regard to murine keratinocytes, tumorigenic lines spontaneously appear in culture



**Figure 1 Differentiation and stratification of murine keratinocytes.**

(A) Calcium induction of Keratin 1. Paired samples of cells were grown to confluence in low calcium and then maintained an additional 48 h in either 0.06 mM calcium (–) or 1.8 mM calcium (+). Protein extracts were separated by PAGE and analyzed by western blotting using anti-K1 antibody (Covance, cat. #PRB-165p-100). Cell lines, with passage number in parentheses, are: (a) BALB/c (17); (b) C57BL/6<sup>SupFG1</sup> (20); (c) SKH (18); (d) 129/SvEvTac<sup>hfe-/-</sup> (22); (e) C57BL/6<sup>hprt-/-</sup> (19). (B) Keratinocyte stratification. 10<sup>6</sup> cells were seeded onto devitalized dermis (Prunieras *et al*, 1983) and cultured submerged for 1 wk and then at the air:liquid interface for 2 wk. (a) C57BL/6; (b) C57BL/6<sup>SupFG1</sup>; (c) C57BL/6<sup>hprt-/-</sup>; (d) SKH. Arrow in (b) indicates keratohyalin granules. Arrow in (d) indicates stratum corneum. H&E.

(Yuspa *et al*, 1980), cells exposed to mutagens readily become polyploid (Kulesz-Martin *et al*, 1983), and cells grown in serum-free medium become polyploid after as few as five passages (Kaighn *et al*, 1988). We had hoped that the method of Hager *et al*, might suppress the tendency to polyploidy.

In our current age of expression profiling and searches for quantitative “downstream” effects of defined mutants, long-term murine keratinocyte cultures should be used with caution.

Victoria L. Bruegel Sanchez, Jing Zhou, Donna LaCivita, and  
Leonard M. Milstone  
Department of Dermatology, Yale University School of Medicine,  
New Haven, Connecticut, USA

---

We are grateful to Barbara Hager and Philip Fleckman for help in mastering their culture method. The studies were supported by NIH-sponsored Tissue Core of the Yale Skin Disease Research Center, P30 AR41942. An abstract of this work was presented at the Society for Investigative Dermatology meeting in 2003.

DOI: 10.1111/j.0022-202X.2004.23218.x

Manuscript received December 23, 2003; revised March 24, 2004; accepted for publication March 30, 2004

Address correspondence to: Leonard M. Milstone, Department of Dermatology, Yale University School of Medicine, New Haven, 06520-8059, USA. Email: leonard.milstone@yale.edu

### References

- Caldelari R, Suter MM, Baumann D, deBruin A, Muller E: Long-term culture of murine epidermal keratinocytes. *J Invest Dermatol* 114:1064–1065, 2000
- Hager B, Bickenbach JR, Fleckman P: Long-term culture of murine epidermal keratinocytes. *J Invest Dermatol* 112:971–976, 1999
- Kaighn ME, Camaliere RF, Bertolero F, Saffiotti U: Spontaneous establishment and characterization of mouse keratinocyte cell lines in serum-free medium. *In Vitro Cell Dev Biol* 24:845–854, 1988
- Kulesz-Martin M, Kilkenny AE, Holbrook KA, Digernes V, Yuspa SH: Properties of carcinogen altered mouse epidermal cells resistant to calcium-induced differentiation. *Carcinogenesis* 4:1367–1377, 1983
- Prunieras M, Regnier M, Woodley D: Methods for cultivation of keratinocytes with an air–liquid interface. *J Invest Dermatol* 81 (1 Suppl.):28s–33s, 1983
- Yuspa SH, Hawley-Nelson P, Koehler B, Stanley JR: A survey of transformation markers in differentiating cell lines in culture. *Cancer Res* 40:4694–4703, 1980